

determine transmission and pathogenicity. As next generation sequencing technologies become mainstream, this should be achievable in the near future.

References

1. Marshall, I.D. (1988) Murray Valley and Kunjin encephalitis. In *The arboviruses: epidemiology and ecology* (Monath, T.P. ed), pp. 151–189, CRC Press, Boca Raton.
2. Selvey, L.A. *et al.* (2014) The changing epidemiology of Murray Valley encephalitis in Australia: the 2011 outbreak and a review of the literature. *PLoS Negl. Trop. Dis.* **8**, e2656. doi:10.1371/journal.pntd.0002656
3. Roche, S.E. *et al.* (2013) Descriptive overview of the 2011 epidemic of arboviral disease in horses in Australia. *Aust. Vet. J.* **91**, 5–13. doi:10.1111/avj.12018
4. French, E.L. *et al.* (1957) Murray Valley encephalitis in New Guinea. I. Isolation of Murray Valley encephalitis virus from the brain of a fatal case of encephalitis occurring in a Papuan native. *Am. J. Trop. Med. Hyg.* **6**, 827–834. doi:10.4269/ajtmh.1957.6.827
5. Lobigs, M. *et al.* (1986) Genetic differentiation of Murray Valley encephalitis virus in Australia and Papua New Guinea. *Aust. J. Exp. Biol. Med. Sci.* **64**, 571–585. doi:10.1038/icb.1986.61
6. Johansen, C.A. *et al.* (2000) Isolation of Japanese encephalitis virus from mosquitoes (Diptera: Culicidae) collected in the Western Province of Papua New Guinea, 1997–1998. *Am. J. Trop. Med. Hyg.* **62**, 631–638. doi:10.4269/ajtmh.2000.62.631
7. Coelen, R.J. and Mackenzie, J.S. (1988) Genetic variation of Murray Valley encephalitis virus. *J. Gen. Virol.* **69**, 1903–1912. doi:10.1099/0022-1317-69-8-1903
8. Coelen, R.J. and Mackenzie, J.S. (1990) The 5'-terminal non-coding region of Murray Valley encephalitis virus RNA is highly conserved. *J. Gen. Virol.* **71**, 241–245. doi:10.1099/0022-1317-71-1-241
9. Johansen, C.A. *et al.* (2007) Genetic and phenotypic differences between isolates of Murray Valley encephalitis virus in Western Australia, 1972–2003. *Virus Genes* **35**, 147–154. doi:10.1007/s11262-007-0091-2
10. Williams, D.T. *et al.* (2015) The molecular epidemiology and evolution of Murray Valley encephalitis virus: recent emergence of distinct sub-lineages of the dominant genotype 1. *PLoS Negl. Trop. Dis.* **9**, e0004240. doi:10.1371/journal.pntd.0004240
11. Mackenzie, J.S. and Broom, A.K. (1999) Ord River irrigation area: the effect of dam construction and irrigation on the incidence of Murray Valley encephalitis virus. In *Water resources: Health, Environment and Development* (Kay, B.H. ed), pp. 108–122, E & FN Spon.
12. Niazi, A. (2014) The role of genetic diversity in the replication, pathogenicity and virulence of Murray Valley encephalitis virus, Curtin University, Perth, Western Australia.
13. Coelen, R. (1988) Phenotypic and genotypic variation of Murray Valley encephalitis virus, Department of Microbiology, University of Western Australia, Perth, Western Australia.
14. Hurrelbrink, R.J. *et al.* (1999) Characterization of infectious Murray Valley encephalitis virus derived from a stably cloned genome-length cDNA. *J. Gen. Virol.* **80**, 3115–3125. doi:10.1099/0022-1317-80-12-3115
15. Mann, R.A. *et al.* (2013) Molecular characterization and phylogenetic analysis of Murray Valley encephalitis virus and West Nile virus (Kunjin subtype) from an arbovirus disease outbreak in horses in Victoria, Australia, in 2011. *J. Vet. Diagn. Invest.* **25**, 35–44. doi:10.1177/1040638712467985
16. Williams, D.T. *et al.* (2014) Complete genome sequences of the prototype isolates of genotypes 2, 3, and 4 of Murray Valley encephalitis virus. *Genome Announc.* **2**. doi:10.1128/genomeA.00581-14
17. Kumar, S. *et al.* (2016) MEGA7: Molecular Evolutionary Genetics Analysis Version 7.0 for Bigger Datasets. *Mol. Biol. Evol.* **33**, 1870–1874. doi:10.1093/molbev/msw054

Biography

Dr David Williams is a Group Leader at the CSIRO Australian Animal Health Laboratory, Geelong, Victoria. His research and scientific interests are in the detection, diagnosis, and epidemiology of emerging and re-emerging viruses that affect humans and animals in Australia and overseas, including mosquito-borne viruses.

Protecting Australia from disease vectors: exotic mosquito management at the border



Angus Sly

Pathway Surveillance and Operational Science Compliance Division
Department of Agriculture and Water Resources
GPO Box 858
Canberra, ACT 2601, Australia
Email: Angus.Sly@agriculture.gov.au



Callum Mack

Office of Health Protection
Australian Government Department of Health
GPO Box 9848
Canberra, ACT 2601, Australia
Email: Callum.Mack@health.gov.au

Mosquitoes, through the diseases they transmit, are considered the deadliest animals in the world¹. While Australia is relatively free of many of the mosquito species capable of transmitting diseases such as dengue, yellow fever,

chikungunya, and Zika virus, Australia is not immune to the risk of these arboviruses becoming endemic through the introduction of exotic mosquito vectors. In 150 separate instances there were 525 individual exotic mosquitoes

detected at the Australian border between 2014 and 2017 (Department of Agriculture and Water Resources, unpublished data). Accordingly, there is a strong focus on surveillance and control activities to prevent exotic mosquito incursions and possible local establishment.

Aedes aegypti (Linnaeus) and *Aedes albopictus* (Skuse) are highly invasive mosquito species responsible for the transmission of diseases significant to public health across many parts of the world. *Aedes albopictus* is not yet established on mainland Australia and *Ae. aegypti* is confined to a limited distribution through eastern Queensland². As a result, diseases such as dengue, chikungunya, Zika virus, and yellow fever are relatively rare or non-existent in Australia. While Queensland continues to experience localised dengue outbreaks stemming from imported cases, the rest of Australia remains free from disease transmission.

The lack of a competent vector dramatically reduces the public health risk posed by arboviral diseases. For example, there were 1740 cases of dengue reported in Western Australia between 2014 and 2017³, predominantly associated with overseas travel. This is the highest recorded of any state or territory during this period, but in the absence of a competent vector no dengue outbreaks have occurred. The establishment of either *Ae. aegypti* or *Ae. albopictus* in this region of Australia would have significant public health consequences.

Australia's island geography aids in preventing the introduction of mosquito vectors, but a reliance on imports and increasing international passenger movements has provided a pathway for invasive mosquitoes via international conveyances and imported cargo. To prevent the establishment of vectors through import pathways and maintain a risk free environment with respect to the previously mentioned arboviruses, the Australian Government employs specific measures at the border, primarily: disinsection of all aircraft entering Australia, targeted inspections of international vessels and cargo, and vector surveillance at first points of entry (air and sea ports).

Exotic mosquito surveillance is legislated under the *Biosecurity Act 2015* and undertaken by the Australian Government Department of Agriculture and Water Resources. In line with the World Health Organization's *International Health Regulations 2005* surveillance is carried out to a minimum distance of 400 metres from facilities that are used for operations involving travellers, conveyances, cargo, and postal articles⁴.

The Department of Agriculture and Water Resources deploys a range of mosquito surveillance traps at first points of entry as part of its mosquito surveillance program. These traps are targeted at collecting different life stages of mosquitoes including Biogents

Sentinel traps and Encephalitis Vector Surveillance CO₂ traps for collecting adults, Sentinel tyre traps for collecting larvae, and ovitraps for collecting mosquito eggs (Figure 1). The vast majority of mosquitoes collected through this program are local species with 117 exotic mosquitoes (37 separate instances) identified from more than half a million mosquitoes collected at Australian first points of entry in 2016 (Department of Agriculture and Water Resources, unpublished data). The type and number of traps deployed at a first point of entry varies according to the size, the volume of arriving international conveyances and cargo, and local environmental conditions at the port. The larger the port or the higher the volume of international traffic arriving at a port, a greater number of traps and trap types are used. Surveillance activities are also escalated at times of heightened international traffic. For example, additional surveillance traps are deployed to cover increased international military arrivals during Australian Defence Force exercises.

The detection of a single exotic mosquito will trigger a rapid response involving Federal, State and, in some jurisdictions, Local Government. A typical response to an exotic mosquito detection will involve knockdown adulticide treatments (Figure 2), residual insecticide harbourage spraying, and treatments of potential mosquito breeding sites. Enhanced surveillance is undertaken to monitor the effectiveness of the treatments applied and to ensure no exotic mosquitoes remain. Site surveys around the detection point are undertaken to ensure that there is no localised breeding of exotic mosquitoes. To assist this cross government response, the Australian Government Department of Health has developed a set of guidelines to support decision making and describe the roles and responsibilities of the key stakeholders involved in the response activities⁵.

Pathway analysis is another important part of the response process. Determining the origin of an exotic mosquito and how it arrived



Figure 1. Cigar shaped *Aedes* mosquito eggs collected from an oviposition trap.



Figure 2. Thermal insecticide fogging treatment being conducted within a baggage unpack area at an International Airport. Image courtesy of Brisbane City Council.

allows for targeted control measures to be implemented. Genome-wide single nucleotide polymorphism (SNP) analysis is used as a population genetics tool to compare genetic similarities between exotic mosquitoes detected at the border and an established reference database in order to determine their origins⁶. This, as well as investigations into insecticide resistance profiles of exotic mosquitoes using genetic markers and biological assays, is helping to better understand new and emerging risk pathways.

To date, pathway controls, early detections of exotic mosquitoes at the border, and rapid response activities have helped prevent exotic mosquito vectors from establishing in Australia, however, exotic mosquitoes continue to probe our borders. Through continued efforts at the border, researchers are provided with more time to develop new and innovative ways to prevent and control these mosquito vectors, and the diseases they transmit ahead of their possible introduction to Australia.

References

1. World Health Organization (2018) Mosquito-borne diseases. http://www.who.int/neglected_diseases/vector_ecology/mosquito-borne-diseases/en/
2. Queensland Health (2015) *Queensland Dengue Management Plan 2015–2020*. <https://www.health.qld.gov.au/publications/clinical-practice/guidelines-procedures/diseases-infection/governance/dengue-mgt-plan.pdf>
3. Australian Government Department of Health (2018) National Notifiable Diseases Surveillance System. <http://www9.health.gov.au/cda/source/cda-index.cfm>
4. World Health Organization (2005) *International Health Regulations*, 3rd edition, World Health Organization, Geneva, 2016.
5. Australian Government Department of Health (2017) Response guide for exotic mosquito detections at Australian first points of entry. <http://www.health.gov.au/internet/main/publishing.nsf/Content/ohp-exotic-mosquitoes>
6. Rašić, G. *et al.* (2014) Genome-wide SNPs lead to strong signals of geographic structure and relatedness patterns in the major arbovirus vector, *Aedes aegypti*. *BMC Genomics* **15**, 275. doi:10.1186/1471-2164-15-275

Biographies

Angus Sly completed a BSc at James Cook University. Mr Sly is an Assistant Director in Pathway Surveillance and Operational Science with the Australian Government Department of Agriculture and Water Resources and is responsible for overseeing the coordination of the department's mosquito surveillance program at Australia's first points of entry.

Callum Mack completed a BSc (Hons) in molecular genetics at the University of New England. A senior policy officer with the Office of Health Protection, part of the Australian Government Department of Health, Mr Mack is responsible for exotic mosquito management policies at Australia's first point of entry.

Short Lab Reports, Book Reviews and Hot Topics of interest to ASM members are invited at any time.