

# Biotechnological perspectives of microorganisms isolated from the Polar Regions



*Viktoria Shcherbakova*

G. K. Skryabin Institute of Biochemistry and Physiology of Microorganisms, Russian Academy of Sciences, Pushchino, Russia  
Tel: +7 9165675019,  
Email: vshakola@gmail.com



*Olga Troshina*

G. K. Skryabin Institute of Biochemistry and Physiology of Microorganisms, Russian Academy of Sciences, Pushchino, Russia  
Tel: +7 9168812865,  
Email: olgtroshina@gmail.com

**Polar permanently frozen grounds cover more than 20% of the earth's surface, and about 60% of the Russian territories are permafrost. In the permafrost environments, the combination of low temperature and poor availability of liquid water make these habitats extremely inhospitable for life. To date, both culture-dependent and culture-independent methods have shown that permafrost is a habitat for microorganisms of all three domains: *Bacteria*, *Archaea* and *Eukarya*. An overview of applying psychrophilic and psychrotolerant bacteria and archaea isolated from Arctic and Antarctic permafrost ecosystems in biotechnological processes of wastewater treatment, production of cold-adapted enzymes, etc. is discussed here. The study of existing collections of microorganisms isolated from permanently cold habitats, improved methods of sampling and enrichment will increase the potential biotechnological applications of permafrost bacteria and archaea producing unique biomolecules.**

Even though microorganisms were found in the permafrost more than 100 years ago, intensive studies of microbial communities in the Arctic and Antarctic began at the end of the last century. Cold-adapted microorganisms do not only survive at low temperatures, but can show significant growth and metabolic activity (respiration and biosynthesis) at temperatures down to  $-20^{\circ}\text{C}$ , and even at  $-39^{\circ}\text{C}$ <sup>1,2</sup>. The number of publications describing new prokaryotic taxa isolated from the polar ecosystems has increased by eight times between 1990 and 2015 (Figure 1).

Cold-adapted microorganisms face diverse challenges inherent to living in cold environments: low enzyme activity and low rates of biochemical reactions, altered transport systems, decreased membrane fluidity, and cold denaturation of proteins<sup>3</sup>. To get over

these challenges, they have developed remarkable adaptations. Psychrophiles produce higher amounts of unsaturated and methyl-branched fatty acids and shorter acyl-chain fatty acids to increase the membrane fluidity<sup>4</sup>. They also produce cold-shock proteins to aid in different cellular processes as well as antifreeze proteins that inhibit ice crystal growth. Furthermore, all components of their cells must be suitably adapted to cold temperatures. The key adaptation to cold is the synthesis of cold-adapted enzymes that show an improved flexibility of enzyme conformation and maintain high specific activities at low temperatures.

The vast majority of current industrial processes are performed under harsh conditions, including extremely high and low temperatures, acidic or basic pH values, and elevated salinity. Standard enzymes have specific requirements for maximal function and perform optimally in narrow ranges of physical and chemical conditions. Cold-adapted enzymes have a combination of specific adaptations in their structural features that give them more flexible structures than mesophilic and thermophilic enzymes. This trait allows for high catalytic activity at low temperatures<sup>5</sup> and




Fig 1. Distribution of the number of new prokaryotic taxa from the Polar regions validated from 1990 to 2015.

thermolability. In recent years, scientific and industrial efforts to discover and develop novel cold-adapted enzymes have increased substantially. The intrinsic characteristics of cold-adapted enzymes, such as high activity at low temperatures and thermolability, are extremely valuable for biotechnological applications in a wide variety of industries from molecular biology to food and beverage preparation<sup>6</sup>. Consequently, psychrophilic enzymes are replacing mesophilic enzymes in many different industrial processes. Due to the limitations of cultivating psychrophiles for production of cold-adapted enzymes, the current strategy is to clone and express genes encoding the desired product in mesophilic hosts prior to the operation in a bioreactor<sup>7</sup>.

Biotechnological applications of psychrophilic and psychrotolerant microorganisms include cold-water detergents, food additives and flavour modifying agents, biosensors, and environmental bioremediation. Cold-active lipases catalyse the hydrolysis of fats and remove fatty acid stains from tissues at low temperatures<sup>8–10</sup>. Recently, high-performance lipase from *Pseudomonas stutzeri* PS59 has been identified and proposed for use. This lipase has its optimal activity at 20°C, pH 8.5, and its activity is not inhibited by various surfactants and oxidants<sup>11</sup>. Despite a number of successes achieved in recent years in the study of cold-active enzymes for the production of detergents, there is a clear need for even more active enzymes from psychroactive microorganisms. Enzymes that can work optimally at low temperatures (15–25°C), but retain high activity over a wide range of temperatures (from 5–60°C), and in the presence of surfactants and at alkaline pH will have future applications in detergent compositions. Enzymes, which possess such universal properties, have already been obtained in the form of recombinant proteins<sup>12</sup>. Proteases are enzymes that catalyse the hydrolysis of peptide bonds in proteins breaking them into shorter fragments. They also play an important role in the washing industry for removal of protein stains. Recent studies have focused on the discovery of new cold-active proteases in psychrophilic microorganisms derived from the Arctic and Antarctic ecosystems<sup>13</sup>.

Psychrophilic and psychroactive microorganisms synthesise various molecules to protect the cell from freezing or to minimise the harmful effects of ice crystal formation<sup>3</sup>. Antifreeze proteins (AFPs) are ice-binding proteins that have the ability to change the crystalline structure of ice and suppress ice growth in two directions. For the first time, the presence of a protein leading to thermal hysteresis (TH) in bacteria was demonstrated by Duman and Olsen<sup>14</sup>, and the *Moraxella* sp. strain became the first bacterial producer of AFPs<sup>15</sup>. At the beginning of studies of the antifreeze phenomenon, the highest values of TH were found in insects (3–6°C) and fish (0.7–1.5°C). AFPs isolated from plants and microorganisms had a significantly smaller TH: from 0.2 to 0.5°C for plants, less than 0.1°C for bacteria. Later, an AFP with high activity was detected in the Antarctic bacterium *Marinomonas*

*primoryensis*<sup>16</sup>, which is able to reduce the freezing point by 2°C. Up until now, the antifreeze activity has been detected in a small number of bacteria, mainly in Antarctic isolates. This fact directly indicates a role for AFPs in bacterial adaptation to cold. To date, antifreeze proteins have been detected in actinobacteria of the genera *Micrococcus* and *Rhodococcus*,  $\gamma$ -proteobacteria of the genera *Colwellia*, *Marinomonas* and *Pseudomonas* and bacteria belonging to the genera *Flavobacterium* and *Chryseobacterium*<sup>17</sup>. Screening of anaerobic prokaryotes in the All-Russian collection of microorganisms (VKM) showed AFP of a new type in the spore-forming bacterium *Clostridium tagluense* strain 121<sup>T</sup> isolated from the Canadian permafrost<sup>18</sup>. Metagenomic analysis of several permafrost Arctic samples of various origins and ages has been carried out<sup>19</sup> and the screening for AFP will be our next step.

Psychrotolerant microorganisms are of great value for bioremediation of hydrocarbon-contaminated soil in Polar Regions, because they can maintain activity under extreme environmental conditions. Several strains of psychrotolerant bacteria have been isolated from the oil-contaminated soil in Antarctica and studied in detail<sup>20,21</sup>. Low-temperature anaerobic digestion (AD) has been successfully applied to treat a vast range of wastewater types, such as phenolic, chlorinated aliphatic, brewery, pharmaceutical and glucose-based wastewaters. The evidence of comparable treatment efficiencies to mesophilic setups has also been recorded, as well as the successful treatment of complex wastewater. Despite successful applications, there is a lack of fundamental knowledge regarding the mechanisms underpinning AD. The future full-scale implementation of AD, and particularly the development of promising new applications, such as low-temperature AD, is severely impaired by this knowledge gap. Methanogenic populations have been the focus of many low-temperature AD studies due to their crucial role in biogas formation and biofilm integrity<sup>22</sup>. As our preliminary tests have shown, permafrost methanogens (Table 1) can be used for the process start-up at moderate and low temperatures. Other possible applications of psychroactive isolates from VKM are also presented in the Table 1.

Nature provides a vast source of biocatalysts. However, the probability of finding the right enzymatic activity for a particular application relies on the available technical capabilities to efficiently assess this large microbial diversity. This capability is mainly mediated by technologies, such as metagenomic screenings, genome mining, and direct enzymatic exploration<sup>23,24</sup>. Metagenomic screenings and genome mining require that the search for a novel enzyme is based on genetic sequence homology to already described enzymes. But the discovery of new enzymes in this way does not always give accurate information, especially for less studied organisms like psychrophiles.

An alternative method is presented in the direct exploration of cold-adapted enzymes and AFPs based on functional screenings of

Table 1. Psychrophilic and psychroactive prokaryotes from Arctic and Antarctic permafrost in VKM.

Species	Source of isolation	Growth temperature (optimum)	Application
<b>Bacteria</b>			
<i>Clostridium algariphilum</i> 14D1 <sup>T</sup> VKM B-2271 <sup>T</sup>	Cryopeg in permafrost, Arctic	0–20°C (5–6°C) <sup>A</sup>	Low temperature waste treatment
' <i>Clostridium frigoriophilum</i> ' 14F <sup>T</sup> VKM B-2368 <sup>T</sup>	Cryopeg in permafrost, Arctic	0–15°C (6°C) <sup>A</sup>	Low temperature waste treatment
<i>Clostridium tagluense</i> 121 VKM B-2271 <sup>T</sup>	Permafrost, Arctic	0–28°C (15°C) <sup>A</sup>	Low temperature waste treatment, AFP producing
' <i>Clostridium deceptionensis</i> ' G4 VKM B-2784 <sup>T</sup>	Volcano ground, Antarctica	0–15°C (6–7°C) <sup>A</sup>	Low temperature waste treatment
<i>Celerinatantimonas yamalensis</i> C7 <sup>T</sup> VKM B-2511 <sup>T</sup>	Cryopeg in permafrost, Arctic	0–34°C (18–22°C) <sup>A</sup>	Nitrogen-fixing at near zero temperature, cold-adapted enzymes
<i>Sphaerochaeta associata</i> GLS2 <sup>T</sup> VKM B-2742 <sup>T</sup>	Permafrost, Arctic	5–37°C (30–34°C)	Low temperature waste treatment
' <i>Psychrobacter murincola</i> ' 1pS VKM B-2269	Cryopeg in permafrost, Arctic	2–37°C (18–20°C) <sup>A</sup>	Cold-adapted enzyme
' <i>Psychrobacter murincola</i> ' 2pS <sup>T</sup> VKM B-2270 <sup>T</sup>	Cryopeg in permafrost, Arctic	2–37°C (16–18°C) <sup>A</sup>	Cold-adapted enzyme
<i>Psychrobacter arcticus</i> VKM B-2377 <sup>T</sup>	Permafrost, Arctic	0–28°C (22°C) <sup>A</sup>	Cold-adapted lipase and esterase
<i>Psychrobacter cryohalolentis</i> VKM B-2378 <sup>T</sup>	Cryopeg in permafrost, Arctic	0–30°C (22°C) <sup>A</sup>	Cold-adapted lipases
' <i>Desulfovibrio algaritolerans</i> ' K3S <sup>T</sup> VKM B-2877 <sup>T</sup>	Cryopeg in permafrost, Arctic	0–36°C (26°C) <sup>A</sup>	Low temperature precipitation of heavy metals
<i>Desulfovibrio arcticus</i> B15 <sup>T</sup> VKM B-2367 <sup>T</sup>	Cryopeg in permafrost, Arctic	0–28°C (24°C) <sup>A</sup>	Low temperature precipitation of heavy metals
<b>Archaea</b>			
' <i>Methanosarcina gilichinskiia</i> ' JL01 <sup>T</sup> VKM B-2370 <sup>T</sup>	Permafrost, Arctic	10–37°C (25–28°C) <sup>A</sup>	Low temperature biogas production
<i>Methanobacterium arcticum</i> M2 <sup>T</sup> VKM B-2371 <sup>T</sup>	Permafrost, Arctic	10–42°C (37°C) <sup>A</sup>	Low temperature biogas production
<i>Methanobacterium veterum</i> MK4 <sup>T</sup> VKM B-2440 <sup>T</sup>	Permafrost, Arctic	5–37°C (30°C) <sup>A</sup>	Low temperature biogas production

<sup>A</sup>Capable of subzero growth.

enzymatic activities in large collections of microorganisms, such as VKM. Studies of microbial communities in cold ecosystems indicate that psychroactive prokaryotes represent a large variety of novel and understudied microorganisms. Research of existing collections of microorganisms isolated from permanently cold habitats, the development of improved methods of sampling and enrichment will increase the potential biotechnological application of permafrost bacteria and archaea producing unique biomolecules.

## References

1. Steven, B. *et al.* (2009) Bacterial and archaeal diversity in permafrost. In *Permafrost Soils. Soil Biology*, vol. 16 (Margesin, R., ed.). pp. 59–72. Springer, Berlin, Heidelberg.
2. Bakermans, C. (2008) Limits for microbial life at subzero temperatures. In *Psychrophiles: From Biodiversity to Biotechnology* (Margesin, R. *et al.*, eds). pp. 17–28. Springer, Berlin, Heidelberg.
3. D'Amico, S. *et al.* (2006) Psychrophilic microorganisms: challenges for life. *EMBO Rep.* 7, 385–389. doi:10.1038/sj.embor.7400662

4. Chintalapati, S. *et al.* (2004) Role of membrane lipid fatty acids in cold adaptation. *Cell Mol. Biol.* **50**, 631–642.
5. Siddiqui, K.S. and Cavicchioli, R. (2006) Cold-adapted enzymes. *Annu. Rev. Biochem.* **75**, 403–433. doi:10.1146/annurev.biochem.75.103004.142723
6. Cavicchioli, R. *et al.* (2011) Biotechnological uses of enzymes from psychrophiles. *Microb. Biotechnol.* **4**, 449–460. doi:10.1111/j.1751-7915.2011.00258.x
7. Karlsson, E.N. *et al.* (1999) Efficient production of truncated thermostable xylanases from *Rhodothermus marinus* in *Escherichia coli* fed-batch cultures. *J. Biosci. Bioeng.* **87**, 598–606. doi:10.1016/S1389-1723(99)80121-2
8. Litantra, R. *et al.* (2013) Expression and biochemical characterization of cold-adapted lipases from Antarctic *Bacillus pumilus* strains. *J. Microbiol. Biotechnol.* **23**, 1221–1228. doi:10.4014/jmb.1305.05006
9. Jiewei, T. *et al.* (2014) Purification and characterization of a cold-adapted lipase from *Oceanobacillus* strain PT-11. *PLoS One* **9**, e101343. doi:10.1371/journal.pone.0101343
10. Ji, X. *et al.* (2015) Purification and characterization of an extracellular cold-adapted alkaline lipase produced by psychrotrophic bacterium *Yersinia enterocolitica* strain KM1. *J. Basic Microbiol.* **55**, 718–728. doi:10.1002/jobm.201400730
11. Li, X.L. *et al.* (2014) A high-detergent-performance, cold-adapted lipase from *Pseudomonas stutzeri* PS59 suitable for detergent formulation. *J. Mol. Catal., B Enzym.* **102**, 16–24. doi:10.1016/j.molcatb.2014.01.006
12. Novototskaya-Vlasova, K. *et al.* (2012) Cloning, purification, and characterization of a cold-adapted esterase produced by *Psychrobacter cryobalolentis* K5<sup>T</sup> from Siberian cryopeg. *FEMS Microbiol. Ecol.* **82**, 367–375. doi:10.1111/j.1574-6941.2012.01385.x
13. Sarmiento, F. *et al.* (2015) Cold and hot extremozymes: industrial relevance and current trends. *Front. Bioeng. Biotechnol.* **3**, 148. doi:10.3389/fbioe.2015.00148
14. Duman, J.G. and Olsen, T.M. (1993) Thermal hysteresis protein activity in bacteria, fungi and phylogenetically diverse plants. *Cryobiology* **30**, 322–328. doi:10.1006/cryo.1993.1031
15. Yamashita, Y. *et al.* (2002) Identification of an antifreeze lipoprotein from *Moraxella* sp. of Antarctic origin. *Biosci. Biotechnol. Biochem.* **66**, 239–247. doi:10.1271/bbb.66.239
16. Gilbert, J.A. *et al.* (2005) A hyperactive, Ca<sup>2+</sup>-dependent antifreeze protein in an Antarctic bacterium. *FEMS Microbiol. Lett.* **245**, 67–72. doi:10.1016/j.femsle.2005.02.022
17. Chattopadhyay, M.K. (2006) Mechanism of bacterial adaptation to low temperature. *J. Biosci.* **31**, 157–165. doi:10.1007/BF02705244
18. Suetin, S.V. *et al.* (2009) *Clostridium tagluense* sp. nov., psychrotolerant anaerobic spore-forming bacterium from Canadian permafrost. *Int. J. Syst. Evol. Microbiol.* **59**, 1421–1426. doi:10.1099/ijs.0.002295-0
19. Rivkina, E. *et al.* (2016) Metagenomic analyses of the late Pleistocene permafrost—additional tools for reconstruction of environmental conditions. *Biogeosciences* **13**, 2207–2219. doi:10.5194/bg-13-2207-2016
20. Aislabie, J. *et al.* (2000) Aromatic hydrocarbon-degrading bacteria from soil near Scott Base, Antarctica. *Polar Biol.* **23**, 183–188. doi:10.1007/s0030000050025
21. Bej, A.K. *et al.* (2000) Cold-tolerant alkane-degrading *Rhodococcus* species from Antarctica. *Polar Biol.* **23**, 100–105. doi:10.1007/s0030000050014
22. Liu, Y. *et al.* (2002) Anaerobic granulation technology of wastewater treatment. *Microb. Biotechnol.* **18**, 99–113. doi:10.1023/A:1014459006210
23. Leis, B. *et al.* (2013) Screening and expression of genes from metagenomes. *Adv. Appl. Microbiol.* **83**, 1–68. doi:10.1016/B978-0-12-407678-5.00001-5
24. Adrio, J.L. and Demain, A.L. (2014) Microbial enzymes: tools for biotechnological processes. *Biomolecules* **4**, 117–139. doi:10.3390/biom4010117

## Biographies

**Dr Viktoria Shcherbakova** is the Head of Laboratory of Anaerobic Microorganisms of VKM Department at G.K. Skryabin Institute of Biochemistry and Physiology of Microorganisms, Russian Academy of Sciences (Russia). She is the Head of Moscow region department of the Russian Interregional Microbiological Society. Her research interests include permafrost microbiology and biology of anaerobic microorganisms.

**Dr Olga Troshina** is a researcher at G.K. Skryabin Institute of Biochemistry and Physiology of Microorganisms. Her research interests focus on microbial ecology, nitrogen and hydrogen metabolism of bacteria, microbial interactions between bacteria and archaea.

## Microbiology Australia


Official Journal of the Australian Society for Microbiology Inc.

### Stay informed

Keep up to date with industry news by subscribing to our email alerts or registering for RSS feeds.  
[www.publish.csiro.au/earlyalert](http://www.publish.csiro.au/earlyalert)



[www.publish.csiro.au/journals](http://www.publish.csiro.au/journals)



The Australian Society  
 for **Microbiology**   
 bringing Microbiologists together